

**Supplemental Material S7.** Significant influence differences based on respondent characteristics.

*This table highlights additional significant ( $p < .001$ ) influence differences between respondents.*

<b>Australia more influenced by this than USA and/or Ireland/UK</b>	<b>Mean Ireland/UK</b>	<b>Mean Australia/NZ</b>	<b>Mean USA</b>
Presence of penetration	2.3	2.7	1.9
Age of the PwD	2.0	2.7	2.3
<b>Ireland/UK less influenced by this than USA or Australia/NZ</b>			
Volume of aspiration	2.6	2.8	2.9
<b>Those working &lt;5 years more influenced by this than those working between 5-10 years and/or &gt;10 years</b>			
	<b>Mean &lt;5 years</b>	<b>Mean Between 5– 10 years</b>	<b>Mean &gt;10 years</b>
Presence of penetration	2.5	2.2	2.0
Presence of cough post swallow	2.9	2.7	2.5
Opinion of colleagues	2.4	2.0	1.8
<b>Hospital-based SLPs more influenced by this than community-based SLPs and or Rehabilitation SLPs</b>			
	<b>Mean Community</b>	<b>Mean Rehabilitation</b>	<b>Mean Hospital</b>
Opinion of colleagues	1.9	2.0	2.4