

Table D. R^2 of individual listener data for both speaker pairs and all morph types in Experiment 2.

	P1					P2				
	FOFF	FOFSL	FOSL	FFSL	Full	FOFF	FOFSL	FOSL	FFSL	Full
f201	.99	.95 ^a	.95	.97	.99	1.00	1.00	.98 ^a	.88	1.00
f202	.92	.98	.87	.83	.98	.98	.93	.95	.74	.97
f203	.95	.99	.73	.39 ^b	.98	.87	.87	.93	.54	.92
f204	.63	.73	.07 ^b	.82	.90	.49 ^b	.65	.28	.75	.74
f206	1.00	.99	.96	.82	.98	.95	1.00 ^a	.99	.39 ^b	.99
f207	.89	.95	.96	.86	.99	.92	.98	.93	.75	.95
f208	1.00 ^a	.99 ^a	.98	.89	.92 ^a	.99	.95 ^a	.94	1.00 ^a	1.00 ^a
f209	.96	.93	.85	.84	.95	.92	.87	.47 ^b	.74	.82
m212	.97	.98	1.00	.96	1.00	.91	.91	.99	.58	.99
m213	.99	.89	.88	.10 ^b	.72	.98	.70	.70	n.c. ^c	.71
m214	.98	.99	.99	.89	.99	.99	.95	.98	.79 ^a	.98
m215	.98	.96 ^a	.98	.81	1.00	.98	.93	.93	.55	.99
m216	1.00	.96	.96	.89	.99	.88	.99	.95	.59	.99
m217	.99	.99	.94	.91	.99	.99	.98	.93	.11	.96
m218	.98	.96	.97	.93	.98	.84	.97	.96	.95	.94
m219	.97	.98	.97	.64	.97	.90	.97	.98	.62	.98
<i>M</i>	.95	.95	.88	.78	.96	.91	.92	.87	.67	.93
<i>SD</i>	.09	.07	.23	.23	.07	.12	.10	.21	.22	.09
<i>MIN</i>	.63	.73	.07	.10	.72	.49	.65	.28	.11	.71
<i>MAX</i>	1.00	.99	1.00	.97	1.00	1.00	1.00	.99	1.00	1.00
<i>N</i>	16	16	16	16	16	16	16	16	15	16

^a) R^2 result from ambiguous fits; ^b) poor fit ($R^2 < .5$) with corresponding PSE and SD value (not listed) replaced by the mean of listener's data of the appropriate condition for the ANOVA. ^c) One fit computation did not converge.