Supplemental material, Searl et al., "Clinical Experiences of People With a Laryngectomy During the SARS COVID-19 Pandemic," *AJSLP*, <u>https://doi.org/10.1044/2021_AJSLP-21-00117</u>

State	n	%	State	n	%
Alaska	1	< 1%	New Hampshire	1	< 1%
Arizona	7	4%	New Jersey	6	4%
California	23	13%	New York	6	4%
Colorado	2	1%	North Carolina	1	< 1%
Connecticut	1	< 1%	North Dakota	1	< 1%
Delaware	1	< 1%	Ohio	10	6%
Florida	10	6%	Oklahoma	3	2%
Georgia	4	2%	Oregon	3	2%
Illinois	5	3%	Pennsylvania	3	2%
Indiana	6	4%	South Carolina	4	2%
Iowa	1	< 1%	South Dakota	1	< 1%
Kansas	1	< 1%	Texas	13	8%
Kentucky	2	1%	Vermont	1	< 1%
Louisiana	3	2%	Virginia	5	3%
Maryland	3	2%	Washington	10	6%
Massachusetts	1	< 1%	West Virginia	4	2%
Michigan	16	9%	Wisconsin	5	3%
Minnesota	4	2%	Not Specified	3	1%
Missouri	2	1%	-		

Supplemental Material S2. Distribution of respondents across states (N = 173). Respondents to the survey, Experiencing the SARS COVID-19 Outbreak as a Person with a Laryngectomy.