

Supplemental Table. Individual descriptive data for participants in each of three groups.

Group	Participant number	Age in years	Years since diagnosis	Hoehn & Yahr stage	Voice and speech severity	Gender
Immediate	02	81	11	2.5	2.5	F
	03	64	2	2	1.5	M
	07	57	5	1	1.6	F
	09	66	1	1.5	2	F
	15	56	3	2	2.1	M
	16	74	8	2	3.0	M
	18	67	1	2	1.9	F
	19	61	4	2	3.6	M
Delayed	01	59	11	2.5	1.5	F
	04	67	5	2.5	1.3	F
	06	54	7	2	2.3	M
	08	74	4	1.25	1.8	F
	10	66	1	3	2.9	M
	12	63	3	2	2.1	M
	13	60	3.5	2	3	M
	20	63	3	1	1	F
Historical	01	67	17	—	4	M
	07	59	8	—	2	M
	08	59	2.5	—	3	F
	12	51	4	—	0	F
	14	76	3	—	4	F
	19	74	17	—	4	F
	23	79	1.5	—	0	M
	27	80	3	—	3	F
	35	61	4	—	3	M
	37	76	20	—	5	M
	38	67	7	—	2	M
	40	75	15	—	1	F
	42	66	8	—	4	M

Note. Immediate = immediate treatment group; Delayed = delayed treatment group; Historical = historical treatment group. Em dashes indicate data not available. Hoehn & Yahr stages are 1–5, with higher stages indicating greater severity. Severity ratings of speech and voice deficits are on a scale of 0–5: 0 = *none*, 1 = *mild*, 3 = *moderate*, 5 = *severe*.